



ARMSTRONG

**NUMERICAL SOLUTION OF A SINGULAR  
INTEGRAL EQUATION ARISING FROM A  
SEQUENTIAL PROBABILITY RATIO TEST**

**Sherwood Samn**

**AEROSPACE MEDICINE DIRECTORATE  
CLINICAL SCIENCES DIVISION  
CLINICAL RESEARCH COORDINATION BRANCH  
2507 Kennedy Circle  
Brooks Air Force Base, TX 78235-5117**

LABORATORY

**March 1995**

**Final Technical Report**



Approved for public release; distribution is unlimited.

19950419 075

DTIC QUALITY INSPECTED 5


**AIR FORCE MATERIEL COMMAND  
BROOKS AIR FORCE BASE, TEXAS**

## NOTICES


When Government drawings, specifications, or other data are used for any purpose other than in connection with a definitely Government-related procurement, the United States Government incurs no responsibility or any obligation whatsoever. The fact that the Government may have formulated or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication, or otherwise in any manner construed, as licensing the holder, or any other person or corporation; or as conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

The Office of Public Affairs has reviewed this technical report, and it is releasable to the National Technical Information Service, where it will be available to the general public, including foreign nationals.

This technical report has been reviewed and is approved for publication.



SHERWOOD SAMN, Ph.D.  
Project Scientist



JOE EDWARD BURTON, Colonel, USAF, MC, CFS  
Chief, Clinical Sciences Division

**REPORT DOCUMENTATION PAGE**Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

<b>1. AGENCY USE ONLY (Leave blank)</b>		<b>2. REPORT DATE</b> March 1995	<b>3. REPORT TYPE AND DATES COVERED</b> Final
<b>4. TITLE AND SUBTITLE</b> Numerical Solution of a Singular Integral Equation Arising From a Sequential Probability Ratio Test		<b>5. FUNDING NUMBERS</b> PE - 62202F PR - 7755 TA - 28 WU - 21	
<b>6. AUTHOR(S)</b>  Sherwood Samn			
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b> Armstrong Laboratory (AFMC) Aerospace Medicine Directorate Clinical Sciences Division, Clinical Research Coordination Branch 2507 Kennedy Circle Brooks Air Force Base, TX 78235-5117		<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>  AL/AO-TR-1995-0020	
<b>9. SPONSORING/MONITORING AGENCY NAMES(S) AND ADDRESS(ES)</b>		<b>10. SPONSORING/MONITORING AGENCY REPORT NUMBER</b>	
<b>11. SUPPLEMENTARY NOTES</b>			
<b>12a. DISTRIBUTION/AVAILABILITY STATEMENT</b>  Approved for public release; distribution is unlimited.		<b>12b. DISTRIBUTION CODE</b>	
<b>13. ABSTRACT (Maximum 200 words)</b>  Integral equations arise naturally in problems in engineering and physics, and their numerical solutions can be obtained directly using popular commercial software packages. The same cannot be said of singular integral equations. This report proposes two numerical methods for solving a singular integral equation that arises in the computation of critical values for a sequential probability ratio test.			
<b>14. SUBJECT TERMS</b> Numerical Solutions Sequential Probability Ratio Test Singular Integral Equation		<b>15. NUMBER OF PAGES</b> 20	
		<b>16. PRICE CODE</b>	
<b>17. SECURITY CLASSIFICATION OF REPORT</b> Unclassified	<b>18. SECURITY CLASSIFICATION OF THIS PAGE</b> Unclassified	<b>19. SECURITY CLASSIFICATION OF ABSTRACT</b> Unclassified	<b>20. LIMITATION OF ABSTRACT</b> UL

## Contents

	Page
STATEMENT OF PROBLEM.....	1
METHOD OF SOLUTION.....	2
RESULTS.....	4
CONCLUSIONS.....	5
References.....	5
APPENDIX.....	7

## List of Figures

<u>Figure</u>		
1	Test Results .....	9
2	Determination of k and -b .....	10
3	q(y) at k* for two sigmas .....	11

<b>Accession For</b>	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
<b>Availability Codes</b>	
Dist	Avail and/or Special
A-1	

# NUMERICAL SOLUTION OF A SINGULAR INTEGRAL EQUATION ARISING FROM A SEQUENTIAL PROBABILITY RATIO TEST

Sherwood Samn  
USAF Armstrong Laboratory

## STATEMENT OF PROBLEM

In a recently derived sequential probability ratio test (SPRT) [1] to discriminate between the two simple hypotheses:  $H_0 : \sigma = \sigma_0$  and  $H_1 : \sigma = \sigma_1$  ( $0 < \sigma_0 < \sigma_1$ ) the relevant probability distribution is given by

$$g_d(z; \sigma) = \begin{cases} \frac{1}{\sqrt{2\pi c}} \frac{1}{\sqrt{z+d}} e^{-\frac{z+d}{2c}}, & \text{for } z > -d \\ 0, & \text{otherwise.} \end{cases}$$

where  $d = \ln \frac{\sigma_1}{\sigma_0} > 0$ , and  $c = c(\sigma) = \frac{\sigma^2}{2} \left( \frac{1}{\sigma_0^2} - \frac{1}{\sigma_1^2} \right) > 0$ . The operating characteristic (OC),  $q$ , satisfies the integral equation [2]

$$q(-b; k, \sigma) = \int_{-\infty}^b g_d(z; \sigma) dz + \int_0^k g_d(z+b; \sigma) q(z; k, \sigma) dz, \quad 0 < -b < k$$

which in this case, with  $y = -b$ , reduces to

$$q(y; k, \sigma) = f_0(y; d, c) + c_1 \int_{k_0(y, d)}^k \frac{1}{\sqrt{z+d-y}} e^{-\frac{z+d-y}{2c}} q(z; k, \sigma) dz, \quad 0 < y < k \quad (1)$$

where  $k_0(y, d) = \max\{0, y - d\}$ ,  $c_1 = c_1(\sigma) = \frac{1}{\sqrt{2\pi c(\sigma)}}$ , and

$$f_0(y; d, c) = \begin{cases} 2\Phi\left(\sqrt{\frac{d-y}{c}}\right) - 1, & \text{for } y < d, \\ 0, & y \geq d. \end{cases}$$

Here  $\Phi$  is the distribution function of the standard normal distribution. The integral is singular for those values of  $y$  satisfying  $k > y \geq d$ .

For given values of  $\alpha, \beta$  ( $0 < \alpha, \beta < 1$ ),  $\sigma_0$ , and  $\sigma_1$  ( $0 < \sigma_0 < \sigma_1$ ), the problem is to find (unique) values  $(k^*, b^*)$  of  $(k, b)$  for which  $q(-b^*; k^*, \sigma_0) = 1 - \alpha$ , and  $q(-b^*; k^*, \sigma_1) = \beta$ .

### METHOD OF SOLUTION

For given values of  $\alpha, \beta$  ( $0 < \alpha, \beta < 1$ ),  $\sigma_0$ , and  $\sigma_1$  ( $0 < \sigma_0 < \sigma_1$ ), let  $b = b_{\alpha, \sigma_0}(k)$  denote the solution of  $q(-b; k, \sigma_0) = 1 - \alpha$ , and  $b = b_{\beta, \sigma_1}(k)$  the solution of  $q(-b; k, \sigma_1) = \beta$ . Then  $k^*$  will merely be the solution of  $b_{\alpha, \sigma_0}(k) = b_{\beta, \sigma_1}(k)$ . Once  $k^*$  is found,  $b^*$  is immediately given by  $b^* = b_{\alpha, \sigma_0}(k^*)$ . Thus, the basic problem is to solve Equation (1) for  $q(y)$  for each given  $k$  and  $\sigma$ .

By defining  $\tilde{q}$  as  $qe^{-\frac{y}{2c}}$ ,  $\tilde{f}_0$  as  $f_0e^{-\frac{y}{2c}}$ , and  $c_2 = c_2(\sigma)$  as  $c_1(\sigma)e^{-\frac{d}{2c(\sigma)}}$ , Equation (1) becomes

$$\tilde{q}(y; k, \sigma) = \tilde{f}_0(y; d, c) + c_2 \int_{k_0(y, d)}^k \frac{\tilde{q}(z; k, \sigma)}{\sqrt{z + d - y}} dz, \quad 0 < y < k \quad (2)$$

With the change of variable  $s = k - z$  in the integral, Equation (2) becomes

$$\tilde{q}(y; k, \sigma) = \tilde{f}_0(y; d, c) + c_2 \int_0^{k - k_0(y, d)} \frac{\tilde{q}(k - s; k, \sigma)}{\sqrt{k + d - y - s}} ds, \quad 0 < y < k \quad (3)$$

Or,

$$\tilde{q}(y; k, \sigma) = \tilde{f}_0(y; d, c) + c_2 \begin{cases} \int_0^k \frac{\tilde{q}(k - s; k, \sigma)}{\sqrt{k + d - y - s}} ds, & 0 < y < \min\{d, k\} \\ \int_0^{k + d - y} \frac{\tilde{q}(k - s; k, \sigma)}{\sqrt{k + d - y - s}} ds, & d \leq y < k \end{cases} \quad (4)$$

This integral equation is solved numerically. We have used two different methods. The first method solves the problem by interpolating the numerator in Equation (4) on each appropriate sub-interval by a quadratic polynomial and then integrating the resulting integrand to obtain a set of weights for the numerical scheme. Details of this method can be found in the Appendix. This method is relatively easy to implement and does not rely on any special software routines other than a solver for a system of linear equations.

The second method makes indirect use of special software routines available commercially. Here the sought function  $\tilde{q}(y)$  is discretized on the interval  $[0, k]$  to yield the approximations  $\hat{q}_n$  of  $\tilde{q}(nh)$ , ( $n = 0, 1, \dots, N$ ), where  $h$  is the (uniform) mesh size and  $k = Nh$ . Because the integral equation (4) is only (weakly) singular when  $d \leq y < k$ , the manner in which it is discretized depends on  $y$ . For  $y < \min\{d, k\}$ , and using Adams method and Backward Differentiation Formulae method [3], one obtains a set of  $N_1 (= \min\{n_d = d/h, N\})$  equations of the form

$$\hat{q}_i = \tilde{f}_0(ih) + c_2 h \left\{ \sum_{j=0}^{p_1-1} \frac{W_{N,j} \hat{q}_{N-j}}{\sqrt{(N-i+n_d-j)h}} + \sum_{j=p_1}^N \frac{\omega_{N-j} \hat{q}_{N-j}}{\sqrt{(N-i+n_d-j)h}} \right\} \quad (5)$$

for  $i = 0, 1, \dots, N_1$ . Here  $W$  and  $\omega$  are the appropriate starting and convolution weights of the method respectively [3], and  $p_1$  is the order of the method which we took to be 4.

For  $y \geq d$  and using a different Backward Differentiation Formulae method [4], one obtains a set of  $N - N_1$  equations of the form

$$\hat{q}_i = \tilde{f}_0(ih) + c_2 \sqrt{\pi h} \left\{ \sum_{j=0}^{2p_2-2} W_{N+n_d-i,j}^a \hat{q}_{N-j} + \sum_{j=2p_2-1}^{N+n_d-i} \omega_{N+n_d-i-j}^a \hat{q}_{N-j} \right\} \quad (6)$$

for  $i = N_1 + 1, \dots, N$ . Here  $W^a$  and  $\omega^a$  are appropriate fractional starting and fractional convolution weights of the method respectively [4], and  $p_2$  is the order of the BDF method which we took to be 4. The weights in both Equation (5) and Equation (6) are obtained from appropriate NAG routines. The two sets of equations constitute a set of  $N + 1$  linear equations in the  $N + 1$  unknowns  $\{\hat{q}_i\}_{i=0}^N$ . It can be solved using any number of standard linear equation solvers. We used one of IMSL's routines [5].

## RESULTS

On the test problem (to be described) as well as the actual problem, the two numerical methods have yielded similar results. Thus we will not distinguish the two in the following discussion.

We tested the numerical method above on a simple problem whose analytical solution is known. The problem consists of solving Equation 1 with a new  $f_0$  given by

$$f_0(y; d, c) = e^{\frac{y}{2c}} \{1 - 2c_2(\sqrt{k + d - y} - \sqrt{\max\{0, y - d\} - (y - d)})\} \quad (7)$$

The analytical solution for this problem is simply

$$q(y; k, \sigma) = e^{\frac{y}{2c}},$$

or,

$$\tilde{q}(y; k, \sigma) \equiv 1$$

The numerical result of solving this simple problem is displayed in Figure 1. Here  $\tilde{q}$  is plotted against  $y$ . In this particular test, the parameters are  $k = 4, \sigma_0 = 1, \sigma_1 = 2$ .  $\tilde{q}(y; k, \sigma)$  was calculated twice, once for  $\sigma = 1$  and once for  $\sigma = 2$ . Both of these are shown in Figure 1. As the analytical solution of  $\tilde{q}$  does not depend on  $\sigma$ , they are exactly the same, and represent faithfully the analytical solution  $\tilde{q} \equiv 1$ . Note, however, that if we had plotted  $q = \tilde{q}e^{\frac{y}{2c}} = e^{\frac{y}{2c}}$  instead, the two would definitely be different, since  $c = c(\sigma)$  is a function of  $\sigma$ .

To illustrate how this method solves the real problem (Equation 1), we pick  $\sigma_0 = 1$  in the null hypothesis  $H_0$  and  $\sigma_1 = 1.2$  in the alternative hypothesis  $H_1$ . Assuming  $\alpha = 0.05$  and  $\beta = 0.1$ , we calculated the curves (in the  $k-y$  plane)  $q(y; k, \sigma_0) = 1 - \alpha$  and  $q(y; k, \sigma_1) = \beta$ . These are displayed in Figure 2. These curves intersect at  $(k^*, y^*) = (4.75, 2.18)$ . Finally, in Figure 3, we plotted  $q(y; k^*, \sigma_0)$  and  $q(y; k^*, \sigma_1)$ . Indeed, when  $y = y^*$ ,  $q(y, k^*, \sigma_0) = 0.95$  and  $q(y, k^*, \sigma_1) = 0.10$ .



## CONCLUSION

In this paper, we have proposed two numerical methods to solve a singular integral equation which arose in the study of a certain Sequential Probability Ratio Test. One method uses an easily-implemented scheme and the other method involves casting the singular portion of the problem into a weakly singular equation of Abel type and making use of proven software for this type of equation ([6] as implemented by NAG [4]) to solve the problem. Both methods have yielded similar results when tested (successfully) on a simple problem and when applied to the real problem.

## References

- [1] Chou, Y. M., Anderson, M., and Samn, S. (In preparation) 1994.
- [2] Ghosh, B. K. Sequential Tests of Statistical Hypotheses. Addison-Wesley Publishing Company, 1970.
- [3] NAG Fortran Library Routine D05BWF.
- [4] NAG Fortran Library Routine D05BYF.
- [5] IMSL MATH/LIBRARY, Routine LSARG.
- [6] Lubich, Ch. Discretized Fractional Calculus. SIAM J. Math. Anal. 17, pp. 704-719, 1986.

## APPENDIX

### A Simple Method for Solving Weakly Singular Integrals

In solving Equation (4), the main difficulty is solving a weakly singular integral of the form

$$\int_a^b \frac{f(x)}{(x-c)^\alpha} dx \quad (\text{A.1})$$

where  $\alpha < 1$  and  $c \leq a < b$ . For the problem in this paper  $\alpha = \frac{1}{2}$ . If we interpolate  $f(x)$  with a (the) polynomial  $P(x) = \sum_{k=0}^n a_k x^k$  of degree  $n$  at the points  $a = x_0 < x_1 < \dots < x_n = b$ , then the integral in (A.1) can be approximated by

$$\int_a^b \frac{\sum_{k=0}^n a_k x^k}{(x-c)^\alpha} dx = \sum_{k=0}^n a_k \int_a^b \frac{x^k}{(x-c)^\alpha} dx = \sum_{k=0}^n a_k w_k = \mathbf{w}^T \mathbf{a}, \quad (\text{A.2})$$

where

$$w_k = \int_a^b \frac{x^k}{(x-c)^\alpha} dx,$$

$\mathbf{w} = (w_0, \dots, w_n)^T$ , and  $\mathbf{a} = (a_0, \dots, a_n)^T$ . It is straight forward to show that

$$w_k = Q_k(b, c) - Q_k(a, c),$$

where for all  $y \leq x$ ,

$$Q_k(x, y) = \sum_{i=0}^k \frac{\binom{k}{i}}{i+1-\alpha} y^{k-i} (x-y)^{i+1-\alpha}.$$

Since the polynomial  $P(x) = \sum_{k=0}^n a_k x^k$  interpolates  $f(x)$  at the  $n+1$  points  $x_0, \dots, x_n$ , each  $a_k$  depends only on  $\{x_i\}_{i=0}^n$  and  $\{f(x_i)\}_{i=0}^n$ . In fact, if  $\mathbf{V}$  is the Vandermonde matrix corresponding to  $\{x_i\}_{i=0}^n$ , then

$$\mathbf{V}\mathbf{a} = \mathbf{f}, \quad (\text{A.3})$$

where  $\mathbf{f} = (f(x_0), \dots, f(x_n))^T$ . Recall the Vandermonde matrix corresponding to  $\{x_i\}_{i=0}^n$  is

$$\mathbf{V} = \begin{pmatrix} 1 & x_0 & x_0^2 & \cdots & x_0^n \\ 1 & x_1 & x_1^2 & \cdots & x_1^n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_n & x_n^2 & \cdots & x_n^n \end{pmatrix}$$

Now by combining Equations (A.2) and (A.3), an approximation to the integral in (A.1) can be written compactly as

$$\mathbf{w}^T \mathbf{V}^{-1} \mathbf{f} \quad (\text{A.4})$$

In solving Equation 4, we have chosen the degree  $n$  of the interpolating polynomial  $P(x)$  to be 3. We could have easily chosen it to be any other larger integer. By repeatedly using Equation (A.4) on successive sets of  $n+1$  mesh points in a chosen partition  $\{0 = x_0, x_1, \dots, x_N = k\}$  of the interval  $[0, k]$  (see Equation (4)), we immediately obtain a system of  $N+1$  linear equations in the  $N+1$  unknowns  $\{f(x_0), f(x_1), \dots, f(x_N)\}$ . This system can now be solved by using any standard system of linear equations solver.

Figure 1. Test Results




Figure 2. Determination of  $k$  and  $-b$




Figure 3.  $q(y)$  at  $k^*$  for two sigmas

